

Carbon sources in fruit carbonate of *Buglossoides arvensis* and consequences for ^{14}C dating

Zamanian K., Pustovoytov K., Kuzyakov Y.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 by the Arizona Board of Regents on behalf of the University of Arizona. Fruit carbonate of *Buglossoides arvensis* (syn. *Lithospermum arvense*) is a valuable dating and paleoenvironmental proxy for late Quaternary deposits and cultural layers because CaCO_3 in fruit is assumed to be accumulated from photosynthetic carbon (C). However, considering the uptake of HCO_3^- by roots from soil solution, the estimated age could be too old depending on the source of HCO_3^- allocated in fruit carbonate. Until now, no studies have assessed the contributions of photosynthetic and soil C to the fruit carbonate. To evaluate this, the allocation of photo-assimilated carbon and root uptake of HCO_3^- was examined by radiocarbon (^{14}C) labeling and tracing. *B. arvensis* was grown in carbonate-free and carbonate-containing soils (sand and loess, respectively), where ^{14}C was provided as (1) $^{14}\text{CO}_2$ in the atmosphere (5 times shoot pulse labeling), or (2) $\text{Na}_2^{14}\text{CO}_3$ in soil solution (root-labeling; 5 times by injecting labeled solution into the soil) during one month of fruit development. Distinctly different patterns of ^{14}C distribution in plant organs after root- and shoot labeling showed the ability of *B. arvensis* to take up HCO_3^- from soil solution. The highest ^{14}C activity from root labeling was recovered in roots, followed by shoots, fruit organics, and fruit carbonate. In contrast, ^{14}C activity after shoot labeling was the highest in shoots, followed by fruit organics, roots and fruit carbonate. Total photo-assimilated C incorporated via shoot labeling in loess-grown plants was 1.51mg lower than in sand, reflecting the presence of dissolved carbonate (i.e. CaCO_3) in loess. Loess carbonate dissolution and root-respired CO_2 in soil solution are both sources of HCO_3^- for root uptake. Considering this dilution effect by carbonates, the total incorporated HCO_3^- comprised 0.15% of C in fruit carbonate after 10 hr of shoot labeling. However, if the incorporated HCO_3^- during 10 hr of shoot labeling is extrapolated for the whole month of fruit development (i.e. 420-hr photoperiod), fruit carbonate in loess-grown plants incorporated approximately 6.3% more HCO_3^- than in sand. Therefore, fruit carbonates from plants grown on calcareous soils may yield overestimated ^{14}C ages around 500 yr because of a few percentage uptake of HCO_3^- by roots. However, the age overestimation because of HCO_3^- uptake becomes insignificant in fruits older than approximately 11,000 yr due to increasing uncertainties in age determination.

<http://dx.doi.org/10.1017/RDC.2016.123>

Keywords

^{14}C labeling, Biogenic carbonate, *Buglossoides arvensis*, *Lithospermum arvense*, Paleoenvironmental proxy, Radiocarbon dating, Reservoir effect

References

- [1] Amiro BD, Ewing LL. 1992. Physiological conditions and uptake of inorganic carbon-14 by plant roots. *Environmental and Experimental Botany* 32: 203-11.
- [2] Aylward GH. 2007. SI chemical data. Milton, Australia: John Wiley & Sons.
- [3] Brix H. 1990. Uptake and photosynthetic utilization of sediment-derived carbon by *Phragmites australis* (Cav.) Trin. ex Steudel. *Aquatic Botany* 38:377-89.
- [4] Cramer MD, Gao ZF, Lips SH. 1999. The influence of dissolved inorganic carbon in the rhizosphere on carbon and nitrogen metabolism in salinity-treated tomato plants. *New Phytologist* 142: 441-50.
- [5] Cramer MD, Lips SH. 1995. Enriched rhizosphere CO concentrations can ameliorate the influence of salinity on hydroponically grown tomato plants. *Plant Physiology* 94:425-32.
- [6] Cramer MD, Richards MB. 1999. The effect of rhizosphere dissolved inorganic carbon on gas exchange characteristics and growth rates of tomato seedlings. *Journal of Experimental Botany* 50:79-87.
- [7] Ford CR, Wurzbarger N, Hendrick RL, Teskey RO. 2007. Soil DIC uptake and fixation in *Pinus taeda* seedlings and its C contribution to plant tissues and ectomycorrhizal fungi. *Tree Physiology* 27:375-83.
- [8] Gocke M, Pustovoytov K, Kuzyakov Y. 2011. Carbonate recrystallization in root-free soil and rhizosphere of *Triticum aestivum* and *Lolium perenne* estimated by C labeling. *Biogeochemistry* 103:209-22.
- [9] Goodfriend GA. 1987. Radiocarbon age anomalies in shell carbonate of land snails from semiarid areas. *Radiocarbon* 29(2):159-67.
- [10] Jahren AH, Amundson R, Kendall C, Wigand P. 2001. Paleoclimatic reconstruction using the correlation in δO of hackberry carbonate and environmental water, North America. *Quaternary Research* 56:252-63.
- [11] Kuzyakov Y, Shevtzova E, Pustovoytov K. 2006. Carbonate re-crystallization in soil revealed by C labeling: Experiment, model and significance for paleo-environmental reconstructions. *Geoderma* 131:45-58.
- [12] Overstreet R, Ruben S, Broyer TC. 1940. The absorption of bicarbonate ion by barley plants as indicated by studies with radioactive carbon. *Proceedings of the National Academy of Sciences USA* 26:688-95.
- [13] Pelkonen P, Vapaavuori EM, Vuorinen H. 1985. HCO uptake through the roots in willow and sunflower and effect of HCO uptake on the productivity of willow cuttings. In: Palz W, Coombs J, Hall DO, editors. *Energy from Biomass*, 3rd E.C. Conference. London: Elsevier. p 417-21.
- [14] Pigati JS, Quade J, Shahanan TM, Haynes CV Jr. 2004. Radiocarbon dating of minute gastropods and new constraints on the timing of late Quaternary spring-discharge deposits in southern Arizona, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology* 204:33-45.
- [15] Pigati JS, Rech JA, Nekola JC. 2010. Radiocarbon dating of small terrestrial gastropod shells in North America. *Quaternary Geochronology* 5:519-32.
- [16] Pustovoytov KE, Riehl S, Mittmann S. 2004. Radiocarbon age of carbonate in fruits of *Lithospermum* from the early Bronze Age settlement of Hirbet ez-Zeraqōn (Jordan). *Vegetation History and Archaeobotany* 13:207-12.
- [17] Pustovoytov K, Riehl S. 2006. Suitability of biogenic carbonate of *Lithospermum* fruits for C dating. *Quaternary Research* 65:508-18.
- [18] Pustovoytov K, Riehl S, Hilger HH, Schumacher E. 2010. Oxygen isotopic composition of fruit carbonate in *Lithospermeae* and its potential for paleoclimate research in the Mediterranean. *Global and Planetary Change* 71:258-68.
- [19] Quade J, Shanying L, Stiner M, Clark AE, Mentzer S. 2014. Radiocarbon dating, mineralogy, and isotopic composition of hackberry endocarps from the Neolithic site of: Asikli Höyük, central Turkey. *Tree-Ring Research* 70:17-25.
- [20] Stolwijk JA, Thimann KV. 1957. On the uptake of carbon dioxide and bicarbonate by roots, and its influence on growth. 1. *Plant Physiology* 32:513-20.
- [21] Viktor A, Cramer MD. 2003. Variation in root-zone CO concentration modifies isotopic fractionation of carbon and nitrogen in tomato seedlings. *New Phytologist* 157:45-54.
- [22] Viktor A, Cramer MD. 2005. The influence of root assimilated inorganic carbon on nitrogen acquisition/assimilation and carbon partitioning. *New Phytologist* 165:157-69.
- [23] Vuorinen AH, Vapaavuori EM, Lapinjoki S. 1989. Time-course of uptake of dissolved inorganic carbon through willow roots in light and in darkness. *Physiol. Plant* 77:33-8.
- [24] Wang Y, Jahren AH, Amundson R. 1997. Potential for C dating of biogenic carbonate in hackberry (*Celtis*) endocarps. *Quaternary Research* 47: 337-43.